

ON THE SIMPLICITY OF LIE ALGEBRA OF LEAVITT PATH ALGEBRA

ADEL ALAHMEDI AND HAMED ALSULAMI

ABSTRACT. For a field F and a row-finite directed graph Γ let $L(\Gamma)$ be the Leavitt path algebra. We find necessary and sufficient conditions for the Lie algebra $[L(\Gamma), L(\Gamma)]$ to be simple.

1. INTRODUCTION.

In [3] G. Abrams and Z. Mesyan found necessary and sufficient conditions for a simple Leavitt path algebra $L(\Gamma)$ to give rise to a simple Lie algebra $[L(\Gamma), L(\Gamma)]$. This result is based on a simple easily checkable criterion for a linear combination of vertices $\sum_i \alpha_i v_i, \alpha_i \in F, v_i \in V$, to lie in $[L(\Gamma), L(\Gamma)]$. In this paper we extend the result of G. Abrams and Z. Mesyan to not necessarily simple algebras and find the necessary and sufficient conditions for a Lie algebra $[L(\Gamma), L(\Gamma)]$ to be simple.

2. DEFINITIONS AND TERMINOLOGY

A (directed) graph $\Gamma = (V, E, s, r)$ consists of two sets V and E that are respectively called vertices and edges, and two maps $s, r : E \rightarrow V$. The vertices $s(e)$ and $r(e)$ are referred to as the source and the range of the edge e , respectively. The graph is called row-finite if for all vertices $v \in V, \text{card}(s^{-1}(v)) < \infty$. A vertex v for which $s^{-1}(v) = \emptyset$ is called a sink. A vertex v such that $r^{-1}(v) = \emptyset$ is called a source. A path $p = e_1 \dots e_n$ in a graph Γ is a sequence of edges $e_1 \dots e_n$ such that $r(e_i) = s(e_{i+1})$ for $i = 1, \dots, n-1$. In this case we say that the path p starts at the vertex $s(e_1)$ and ends at the vertex $r(e_n)$. If $s(e_1) = r(e_n)$, then the path is closed. If $p = e_1 \dots e_n$ is a closed path and the vertices $s(e_1), \dots, s(e_n)$ are distinct, then the subgraph $(s(e_1), \dots, s(e_n); e_1, \dots, e_n)$ of the graph Γ is called a cycle. A cycle of length 1 is called a loop.

Definition 1. Let W be a subset of V . We say that

- W is hereditary if $v \in W$ implies $w \in W$ for every vertex w connects to v .
- W is saturated if $\{r(e) : s(e) = v\} \subseteq W$ implies that $v \in W$, for every non-sink vertex $v \in V$.

Definition 2. We call an edge $e \in E$ a *fiber* if $s(e)$ is source, $r(e)$ is sink and $E(V, r(e)) = \{e\}$.

Key words and phrases. Leavitt Path Algebra. Cuntz-Krieger C^* -Algebras. Simple Lie Algebra.

Definition 3. We call a vertex v in a connected graph $\Gamma(V, E)$ a *balloon* over a nonempty subset W of V if (i) $v \notin W$, (ii) there is a loop $C \in E(v, v)$, (iii) $E(v, W) \neq \emptyset$, (iv) $E(v, V) = \{C\} \cup E(v, W)$, and (v) $E(V, v) = \{C\}$.

Let Γ be a row-finite graph and let F be a field. The Leavitt path F -algebra $L(\Gamma)$ is the F -algebra presented by the set of generators $\{v | v \in V\}$, $\{e, e^* | e \in E\}$ and the set of relators (1) $v_i v_j = \delta_{v_i, v_j} v_i$ for all $v_i, v_j \in V$; (2) $s(e)e = er(e) = e$, $r(e)e^* = e^*s(e) = e^*$ for all $e \in E$; (3) $e^*f = \delta_{e,f}r(e)$, for all $e, f \in E$; (4) $v = \sum_{s(e)=v} ee^*$, for an arbitrary vertex v which is not a sink. The mapping which sends v to v for $v \in V$, e to e^* and e^* to e for $e \in E$, extends to an involution of the algebra $L(\Gamma)$. If $p = e_1 \dots e_n$ is a path, then $p^* = e_n^* \dots e_1^*$. In what follows we consider only row-finite directed graphs. We call a graph Γ simple if the Leavitt path algebra $L(\Gamma)$ is simple. The conditions for a graph to be simple are given in [1].

Let A be an associative F -algebra. For elements $a, b \in A$, let $[a, b] = ab - ba$ be their the commutator. Then $A^{(-)} = (A, [,]) is a Lie algebra. If A is an associative algebra and S is a subset of A , we will denote the ideal of A generated by S as $id_A(S)$.$


3. LIE ALGEBRA OF LEAVITT PATH ALGEBRA


We start with theorem by G. Abrams and Z. Mesyan in [3].

Theorem 1. ([3]) *Let $\Gamma(V, E)$ be a directed graph. Let $L(\Gamma)$ be a simple algebra.*

- (i) *If V is infinite then the Lie algebra $[L(\Gamma), L(\Gamma)]$ is simple;*
- (ii) *If V is finite, then $[L(\Gamma), L(\Gamma)]$ is simple if and only if $1_{L(\Gamma)} = \sum_{v \in V} v \notin [L(\Gamma), L(\Gamma)]$.*

There exist however non-simple Leavitt path algebras having the Lie algebra $[L(\Gamma), L(\Gamma)]$ simple.




Example 1. Let $\Gamma =$ . The Lie algebra $[L(\Gamma), L(\Gamma)]$ is isomorphic to the Lie algebra of infinite finitary matrices over the Leavitt algebra $L(2)$ and therefore is simple.

The following theorem gives a classification of directed graph having $[L(\Gamma), L(\Gamma)]$ simple.

Theorem 2. *Let $\Gamma(V, E)$ be a directed row-finite graph. The Lie algebra $[L(\Gamma), L(\Gamma)]$ is simple if and only if either $L(\Gamma)$ is simple- this case is covered by Theorem1 - or Γ contains a simple subgraph W such that every point $v \in V \setminus W$ is a balloon over W , and $\sum_{w \in r(E(v, W))} w \in [L(W), L(W)]$.*

We will prove the theorem by proving a series of lemmas. The first lemma is due to G. Abrams and Z. Mesyan, [3]. We will state it without proof.

Lemma 1. ([3]) Let $\Gamma(V, E)$ be a directed graph. Then $[L(\Gamma), L(\Gamma)] = (0)$ if and

only if Γ is a disjoint union of \bullet ,  (vertices and loops).


Lemma 2. Let $\Gamma(V, E)$ be a row-finite graph. If the Lie algebra $[L(\Gamma), L(\Gamma)]$ is nonzero simple, then every cycle has an exit.

Proof. Let C be a no exist cycle of Γ of length d . Then $L(C) \cong M_d(F[t, t^{-1}])$. Let a be the sum of all vertices on the cycle C . The element a is the identity of $L(C)$ and $L(C) = aL(\Gamma)a$. Consider the ideal $J_n = (1-t)^n F[t, t^{-1}]$ of $F[t, t^{-1}]$. Now, if $d \geq 2$, then $[M_d(J_n), M_d(J_n)] \neq (0)$ for all $n \geq 1$, see [5]. Let $I_n = id_{L(\Gamma)}(M_d(J_n))$. Then $[I_n, I_n] \triangleleft [L(\Gamma), L(\Gamma)]$ and because of simplicity of $[L(\Gamma), L(\Gamma)]$ we have $[L(\Gamma), L(\Gamma)] = [I_n, I_n] \subseteq I_n$. Hence $[L(\Gamma), L(\Gamma)] \cap L(C) \subseteq I_n \cap L(C) = M_d(J_n)$. Since $\cap_n J_n = (0)$, it follows that $[L(\Gamma), L(\Gamma)] \cap L(C) = (0)$, but $(0) \neq [M_d(F[t, t^{-1}]), M_d(F[t, t^{-1}])] \subseteq [L(\Gamma), L(\Gamma)] \cap L(C)$. A contradiction. Hence $d = 1$. Thus C is a loop. Since C has no exit and can not be isolated there exist an edge $e \in E$, such that $s(e) \notin V(C) = \{v\}$. Let $J_n = (v - C)^n L(C)$, $I_n = id_{L(\Gamma)}(J_n)$, $vI_nv \subseteq J_n$. Now, $[eJ_n, J_n] = eJ_n \neq (0)$. Hence $[I_n, I_n] \neq (0)$, $[L(\Gamma), L(\Gamma)] = [I_n, I_n] \subseteq I_n$ and therefore $v[L(\Gamma), L(\Gamma)]v \subseteq J_n$. Since $\cap_n J_n = (0)$ it follows that $v[L(\Gamma), L(\Gamma)]v = (0)$, but $[e^*, e] = v - ee^*$, and $v[e^*, e]v = v \neq 0$. A contradiction. \square

The algebra $L(\Gamma)$ is graded: $dg(v) = 0$, $dg(e) = 1$, $dg(e^*) = -1$ for all $v \in V, e \in E$. In [9] it is shown that every graded ideal I of $L(\Gamma)$ is generated (as an ideal) by $I \cap V$. Thus there is a one-to-one correspondence between graded ideals and hereditary saturated subsets of V .

Lemma 3. Let $\Gamma(V, E)$ be a row-finite graph. Let W be nonempty hereditary and saturated subset of V . Let $I = id_{L(\Gamma)}(W)$. If $[L(\Gamma), L(\Gamma)]$ is nonzero simple, then $[I, I] \neq (0)$.


Proof. If $[I, I] = (0)$, then, in particular, $[L(W), L(W)] = (0)$, W is a disjoint

union of \bullet , and . This implies that for every vertex $w \in W$ there exist an edge $e \in E$ such that $r(e) = w$, $s(e) \notin W$ otherwise w is isolated in Γ . Now, $e, e^* \in I$ and $[e, e^*] \neq 0$. Lemma is proved. \square

Lemma 4. Let $\Gamma(V, E)$ be a row-finite graph. If $[L(\Gamma), L(\Gamma)]$ is nonzero simple, then there exists a minimal hereditary saturated subset in V .

Proof. We need to show that the intersection of all nonzero graded ideals in $L(\Gamma)$ is nonzero. If I is a nonzero graded ideal of $L(\Gamma)$ then by Lemma 3 $[I, I] \neq (0)$. Since $[L(\Gamma), L(\Gamma)]$ is simple, then $[L(\Gamma), L(\Gamma)] = [I, I]$ and therefore $[L(\Gamma), L(\Gamma)]$ lies in the intersection of all nonzero graded ideals of $L(\Gamma)$. \square



Let $\Gamma(V, E)$ be a row-finite graph. Suppose $[L(\Gamma), L(\Gamma)]$ is nonzero simple. Let W be a minimal hereditary saturated subset in V . Let $I = id_{L(\Gamma)}(W)$, $\Gamma' = (V \setminus W, E \setminus E(V, W))$. We assume that $W \neq V$, that is $L(\Gamma)$ is not simple. Since $L(\Gamma') \cong L(\Gamma)/I$ and $[L(\Gamma), L(\Gamma)] \subseteq I$ it follows that $[L(\Gamma'), L(\Gamma')] = (0)$. By Lemma 1 Γ' is

a disjoint union of \bullet , and .

Lemma 5. Γ' does not have components \bullet .

Proof. Let a vertex $v \in V \setminus W$ be isolated in Γ' . Then $E(V \setminus W, v) = \emptyset$. Since W is hereditary and $v \notin W$ we conclude that $E(V, v) = \emptyset$. Since v can not be isolated in Γ it can not be a sink, $E(v, V) \neq \emptyset$. But $E(v, V \setminus W) = \emptyset$, hence all descendants of v lie in W . Since W is saturated we conclude that $v \in W$, a contradiction. \square

Lemma 6. Every vertex $v \in V \setminus W$ is a balloon over W .

Proof. By what we have shown Γ' is a disjoint union of loops . It is easy to see that $E(V, v) = \{c\}$ and $E(v, V \setminus W) = \{c\}$. If $E(v, W) = \emptyset$ then the loop  is isolated in Γ . Hence $E(v, W) \neq \emptyset$. Thus v is a balloon over W . \square

Let S_0 be the span of all elements pp^* , where p is a path on Γ including pathes of length zero (that is vertices). Let S_1 be the span of all elements pq^* , where p, q are pathes on Γ , $r(p) = r(q)$, $p \neq q$. It follows from the description of a Groebner - Shirshov basis of $L(\Gamma)$ [4] that $L(\Gamma) = S_0 + S_1$ is a direct sum of vector spaces. Let M be the semigroup generated by $V \cup E \cup E^*$. It is easy to see that (i) $M = (M \cap S_0) \cup (M \cap S_1)$, (ii) for arbitrary elements $a, b \in M$ if $0 \neq ab \in S_i$, then $ba \in S_i$ or $ba = 0$, for $i = 0, 1$.

Lemma 7. $[I, I] \cap S_0 = \text{span}\{[p, p^*] \mid p \text{ is a path on } \Gamma, r(p) \in W\}$.

Proof. The ideal I is spanned by elements pq^* ; p, q are paths, $r(p) = r(q) \in W$. Consider two such elements $p_1q_1^*$ and $p_2q_2^*$, $0 \neq p_1q_1^*p_2q_2^* \in S_0$. Since $q_1^*p_2 \neq 0$ it follows that $p_2 = q_1u$ or $q_1 = p_2u$, where u is a path on Γ . Consider the first case, $p_2 = q_1u$. Then $p_1q_1^*p_2q_2^* = p_1uq_2^*$. Since this element lies in S_0 we conclude that $q_2 = p_1u$. Now, $p_2q_2^*p_1q_1^* = q_1uu^*p_1^*p_1q_1^* = (q_1u)(q_1u)^*$ and therefore $[p_1q_1^*, p_2q_2^*] = (p_1u)(p_1u)^* - (q_1u)(q_1u)^* = [p_1u, (p_1u)^*] - [q_1u, (q_1u)^*]$. Remember that $r(u) = r(q_2) \in W$. Let $q_1 = p_2u$. Then $p_1q_1^*p_2q_2^* = p_1u^*p_2^*p_2q_2^* = p_1(q_2u)^*$. Again $p_1q_1^*p_2q_2^* \in S_0$ implies $p_1 = q_2u$. Now, $p_2q_2^*p_1q_1^* = p_2q_2^*q_2uu^*p_2^* = (p_2u)(p_2u)^*$. Therefore, $[p_1q_1^*, p_2q_2^*] = (q_2u)(q_2u)^* - (p_2u)(p_2u)^* = [q_2u, (q_2u)^*] - [p_2u, (p_2u)^*]$ and $r(u) = r(p_1) \in W$. \square

Let $v \in V \setminus W$, $E(v, W) = \{e_1, \dots, e_n\}$, $r(e_i) = w_i$ for $1 \leq i \leq n$. Let $w = \sum_{i=1}^n w_i$.

Lemma 8. $w \in [L(W), L(W)]$.

Proof. Since v is a balloon over W , let c be the loop from $E(v, v)$, we have $v = cc^* + \sum_{i=1}^n e_i e_i^*$. Hence $c^*c - cc^* = v - (v - \sum_{i=1}^n e_i e_i^*) = \sum_{i=1}^n e_i e_i^* = \sum_{i=1}^n [e_i, e_i^*] + \sum_{i=1}^n e_i^* e_i = \sum_{i=1}^n [e_i, e_i^*] + w$. Thus $w = c^*c - cc^* - \sum_{i=1}^n [e_i, e_i^*] = [c^*, c] - \sum_{i=1}^n [e_i, e_i^*] \in [L(\Gamma), L(\Gamma)] = [I, I]$. Hence $w \in [I, I] \cap S_0$. By Lemma 7 $w = \sum_i \alpha_i [p_i, p_i^*]$, $\alpha_i \in F$, $r(p_i) \in W$.

We will distinguish between pathes that start with an edge from $E(V \setminus W, W)$ and pathes that lie entirely on W , $w = \sum_i \alpha_{e,i} [ep_{e,i}, p_{e,i}^* e^*] + \sum \beta [q, q^*]$, where e runs over

$E(V \setminus W, W)$, $\alpha_{e,i} \in F$, $p_{e,i}$ and q are paths on W . We have, $w = \sum_i \alpha_{e,i}(ep_{e,i}p_{e,i}^*e^* - r(p_{e,i})) + \sum \beta[q, q^*]$. Fix $e \in E(V \setminus W, W)$. From the description of the basis of $L(\Gamma)$ in [4] it follows that $\sum_i \alpha_{e,i}ep_{e,i}p_{e,i}^*e^* = 0$ and therefore $\sum_i \alpha_{e,i}p_{e,i}p_{e,i}^*e^* = 0$. Now $\sum_i \alpha_{e,i}(ep_{e,i}p_{e,i}^*e^* - r(p_{e,i})) = \sum_i \alpha_{e,i}[p_{e,i}, p_{e,i}^*] \in [L(W), L(W)]$. Hence $w = \sum \alpha_{e,i}[p_{e,i}, p_{e,i}^*] + \sum \beta[q, q^*] \in [L(W), L(W)]$. \square

We proved Theorem 2 in one direction.

4. SIMPLICITY OF THE LIE ALGEBRA OF LEAVITT PATH ALGEBRA

Let $\Gamma(V, E)$ be a graph. Suppose that $W \subsetneq V$ is a simple subgraph, every vertex $v \in V \setminus W$ is a balloon over W and $\sum_{w \in r(E(v, W))} w$ lies in $[L(W), L(W)]$. We will show that the algebra $[L(\Gamma), L(\Gamma)]$ is simple. As above, denote $I = id_{L(\Gamma)}(W)$. The following lemma was proved in [5].

Lemma 9. *I is a simple algebra.*

Lemma 10. *Let A be an arbitrary simple algebra with two orthogonal idempotents e_1, e_2 . Then $A = [A, A] + e_i A e_i$, $i = 1, 2$.*

Proof. We have $A = Ae_1A$. For arbitrary elements $a, b \in A$, $ae_1b = [a, e_1b] + e_1ba$. Similarly, $A = Ae_2A$. For arbitrary elements $a, b \in A$, we have $e_1ae_2b = [e_1ae_2, e_2b] + e_2be_1ae_2$. We proved that $A = [A, A] + e_2Ae_2$. The equality $A = [A, A] + e_1Ae_1$ is proved similarly. \square

Lemma 11. $[L(\Gamma), L(\Gamma)] = [I, I]$.

Proof. We have $L(\Gamma) = I + \text{span}\{c_v^n | n \geq 0, v \in V \setminus W\} + \text{span}\{(c_v^*)^n | n \geq 1, v \in V \setminus W\}$. Let $w \in W$. Then, by Lemma 10, $I = [I, I] + wIw$. Hence $[c_v^n, I] = [c_v^n, [I, I] + wIw] = [c_v^n, [I, I]] \subseteq [I, I]$. Similarly, $[(c_v^*)^n, I] \subseteq [I, I]$. It remains to show that $[c_v^n, (c_v^*)^m] \in [I, I]$. Let $c = c_v$. Suppose at first that $m > n$. Then

$$\begin{aligned} [c^n, (c^*)^m] &= c^n(c^*)^m - (c^*)^{m-n} \\ &= c^{n-1}(cc^*)(c^*)^{m-1} - (c^*)^{m-n} \\ &= c^{n-1}(v - \sum e_i e_i^*)(c^*)^{m-1} - (c^*)^{m-n} \\ &= (c^{n-1}(c^*)^{m-1} - (c^*)^{m-n}) - c^{n-1} \sum e_i e_i^* (c^*)^{m-1}. \end{aligned}$$

The first summand $c^{n-1}(c^*)^{m-1} - (c^*)^{m-n} = [c^{n-1}, (c^*)^{m-1}]$ and we can apply the induction assumption. Furthermore, $c^{n-1}e_i e_i^* (c^*)^{m-1} = [c^{n-1}e_i, e_i^* (c^*)^{m-1}] + e_i^* (c^*)^{m-1} c^{n-1} e_i = [c^{n-1}e_i, e_i^* (c^*)^{m-1}]$, since $e_i^* (c^*)^{m-1} c^{n-1} e_i = e_i^* (c^*)^{m-n} e_i = 0$. Now, let $n > m$. Then

$$\begin{aligned} [c^n, (c^*)^m] &= c^n(c^*)^m - c^{n-m} \\ &= c^{n-1}(v - \sum e_i e_i^*)(c^*)^{m-1} - c^{n-m} \\ &= [c^{n-1}, (c^*)^{m-1}] - c^{n-1} \sum e_i e_i^* (c^*)^{m-1}. \end{aligned}$$

As above,

$$\begin{aligned} c^{n-1}e_i e_i^* (c^*)^{m-1} &= [c^{n-1}e_i, e_i^* (c^*)^{m-1}] + e_i^* (c^*)^{m-1} c^{n-1}e_i \\ &= [c^{n-1}e_i, e_i^* (c^*)^{m-1}] + e_i^* c^{n-m}e_i = [c^{n-1}e_i, e_i^* (c^*)^{m-1}]. \end{aligned}$$

Finally, let $n = m$. As above we conclude that

$$[c^n, (c^*)^n] = [c^{n-1}, (c^*)^{n-1}] - c^{n-1} \sum e_i e_i^* (c^*)^{n-1},$$

$$\sum c^{n-1}e_i e_i^* (c^*)^{n-1} = \sum [c^{n-1}e_i, e_i^* (c^*)^{n-1}] + \sum e_i^* e_i \in [I, I] \text{ by our assumption.}$$

Lemma is proved. \square

Lemma 12. *The algebra $[I, I]$ has zero center.*

Proof. I. Herstein [7] proved that in a simple associative algebra A of dimension bigger than 4 over its center, $[A, A]$ generates A . Hence an elements from I , that commutes with $[I, I]$, lies in the center of I . An arbitrary element from I looks

$$\text{as } z = a_0 + \sum_{e \in E(v_i, W)} ea_e + \sum_{e \in E(v_i, W)} b_e^* e^* + \sum_{\substack{e \in E(v_i, W) \\ f \in E(v_j, W)}} ea_{e,f} f^*, \quad a_0, a_e, b_e, a_{e,f} \in$$

$L(W)$. Suppose that z lies in the center of I . Commuting z with idempotents $w \in W$, $ee^*, e \in E(v_i, W)$ we see that $z = a_0 + \sum_{e \in E(v_i, W)} ea_e e^*$. This implies that a_0

lies in the center of W . Therefore by [2], $|W| < \infty$ and $a_0 = \alpha \sum_{w \in W} w$, $\alpha \in F$.

Multiplying z on the left by e^* and on the right by e , $e \in E(v_i, W)$, we get $r(e)z = a_e = \alpha r(e)$. We proved that $z = \alpha(\sum_{w \in W} w + \sum_{e \in E(v_i, W)} ee^*)$. Now choose

a vertex $v_i \in V \setminus W$ and an edge $f \in E(v_i, W)$. We have $zc_i f = 0$, whereas $c_i f z = \alpha c_i f$. Hence $\alpha = 0$. Lemma is proved. \square

Now it remans to refer to Herstein's theorem about simplicity of $[I, I]/\text{center}$, see [8]. Hence $[I, I]$ is simple and therefore $[L(\Gamma), L(\Gamma)]$ is simple.

ACKNOWLEDGEMENT

The authors would like to thank professor Efim Zelmanov for his constant advise and valuable help during the preparation of this work. The authors would also like to express their appreciation to professor S. K. Jain for carefully reading the manuscript and for offering his comments. This paper was funded by King Abdulaziz University, under grant No. (7-130/1433 HiCi). The authors, therefore, acknowledge technical and financial support of KAU.

REFERENCES

1. G. Abrams, G. Aranda Pino, *The Leavitt path algebra of a graph*, J. Algebra 293 (2005), 319-334
2. G. Aranda Pino, K. Crow *The center of a Leavitt path algebra*, Rev. Mat. Iberoamericana Volume 27, Number 2 (2011), 621-644
3. G. Abrams, Z. Mesyan, *Simple Lie algebra arising from Leavitt path algebra*, Journal of pure and applied algebra, 216(2012), 2303-2313.
4. A. Alahmadi, H. Alsulami, S.K. Jain, E. Zelmanov, *Leavitt path algebras of finite Gelfand Kirillov dimension*, Journal of Algebra and Its Applications, 171(2012)

5. A. Alahmadi, H. Alsulami, *Simplicity of Lie algebra of skew elements of Leavitt path algebra*, submitted
6. P. Colak, *Two-sided ideals in Leavitt path algebras*, Journal of Algebra and Its Applications 10-5 (2011)
7. I.N.Herstein, *Topics in Ring Theory*, Mathematics Lecture Notes, University of Chicago, 1965
8. I.N.Herstein, *Rings with Involution*, Mathematics Lecture Notes, University of Chicago, 1976.
9. Mark Tomforde, *Uniqueness theorems and ideal structure for Leavitt path algebras*, Journal of Algebra 318 (2007), 270299

DEPARTMENT OF MATHEMATICS, KING ABDULAZIZ UNIVERSITY, P.O.Box 80203, JEDDAH, 21589, SAUDI ARABIA

Current address: Department of Mathematics, King Abdulaziz University, P.O.Box 80203, Jeddah, 21589, Saudi Arabia

E-mail address: adelnife2@yahoo.com

DEPARTMENT OF MATHEMATICS, KING ABDULAZIZ UNIVERSITY, P.O.Box 80203, JEDDAH, 21589, SAUDI ARABIA

E-mail address: hhaalsalmi@kau.edu.sa